

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 48/00, C12N 5/00, 5/10	A1	(11) International Publication Number: WO 00/62818 (43) International Publication Date: 26 October 2000 (26.10.00)
(21) International Application Number: PCT/US00/08793 (22) International Filing Date: 3 April 2000 (03.04.00) (30) Priority Data: 09/294,494 20 April 1999 (20.04.99) US (71)(72) Applicant and Inventor: EDELSON, Richard, Leslie [US/US]; 76 Coleytown Road, Westport, CN 06880 (US). (74) Agent: MOORE, Steven, J.; Cummings & Lockwood, 700 State Street, P.O. Box 1960, New Haven, CT 06509-1960 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: DIFFERENTIATION OF MONOCYTES INTO FUNCTIONAL DENDRITIC CELLS (57) Abstract A method for inducing differentiation of monocytes contained in an extracorporeal quantity of a subject's blood into functional dendritic antigen presenting cells is provided. The monocytes are first treated by exposure to physical perturbation, irradiation in the presence of a photoactivatable agent capable of forming photoadducts with cellular DNA components, and/or treatment with a DNA binding agent. The treated monocytes are then incubated for a period of time sufficient to maximize the number of functional dendritic cells in the treated cell population. Functional dendritic cells generated from induced monocytes are incubated together with disease effector agents to enhance the presentation of at least one disease-causing antigen expressed by the disease effector agents. Compositions including dendritic cells derived from induced monocytes and compositions including such dendritic cells incubated with disease effector agents are also provided.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LJ	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

- 1 -

DIFFERENTIATION OF MONOCYTES INTO FUNCTIONAL DENDRITIC CELLS

Field of the Invention

The present invention relates to *in vivo* methods for inducing the differentiation of monocytes into functional dendritic antigen presenting cells and, more particularly, to extracorporeal methods for treating and incubating monocytes to induce such differentiation. The present invention further provides immunotherapeutic compositions including these dendritic cells. In particular, the present invention provides immunotherapeutic compositions comprising apoptotic or inactivated disease effector agents and functional dendritic cells derived from induced monocytes.

Background of the Invention

The use of dendritic cells in cancer immunotherapy is presently an area of significant clinical inquiry. Dendritic cells are highly effective in presenting antigens to responding T-cells; however, dendritic cells normally constitute less than one percent of blood mononuclear leukocytes. Accordingly, a number of *in vitro* methods have been developed to expand populations of dendritic cells to augment anti-cancer immunity. By exposing increased numbers of dendritic cells to antigens on tumor or other disease-causing cells, followed by reintroduction of the antigen-loaded dendritic cells to the patient, presentation of these antigens to responding T-cells can be enhanced significantly.

For example, culturing blood mononuclear leukocytes for eight days in the presence of granulocyte-monocyte colony stimulating factor (GM-CSF) and interleukin-4 (IL-4) produces large numbers of dendritic cells. These cells can then be externally loaded with tumor-derived peptide antigens for presentation to T-cells. Alternatively, the dendritic cells can be transduced to produce and present these antigens themselves. Expanding populations of dendritic cells transduced to produce and secrete cytokines which recruit and activate other mononuclear leukocytes,

- 2 -

including T-cells, may be an even more effective method of generating anti-tumor immune responses.

Transducing cultivated dendritic cells to produce a particular generic tumor antigen and/or additional cytokines is labor intensive and expensive. More importantly, this procedure likely fails to produce and present those multiple tumor antigens that may be most relevant to the individual's own cancer. Several approaches have been proposed to overcome this problem. Hybridization of cultivated autologous dendritic cells with tumor cells would produce tetraploid cells capable of processing and presenting multiple unknown tumor antigens. In a second proposed approach, acid elution of Class I and Class II major histocompatibility complexes (MHC) from the surface of malignant cells would liberate a broad spectrum of tumor-derived peptides. These liberated peptides could then be externally loaded onto MHC complexes of autologous cultivated dendritic cells.

Conventional photopheresis is a method of vaccinating patients against leukemic lymphocytes, even when the distinctive tumor antigen(s) is not known. In this method, malignant cells are exposed to photo-activated 8-methoxypsoralen (8-MOP) which enhances cell surface display of Class I MHC-associated tumor antigens. After intravenous return of these altered malignant lymphocytes to the original patient, a potent anti-tumor response may be generated in about 25% of the patients, leading to diminution of the malignant cell population and occasionally long-standing remissions. Experimental studies in mice, in which autologous dendritic cells are first grown in tissue culture and then admixed with the 8-MOP-treated tumor cells, appears to increase the efficacy of conventional photopheresis. In this experimental protocol, tumorigenic mouse T-cells are rendered apoptotic by photopheresis using 8-MOP and exposure to ultraviolet (UV) energy. Following this chemical alteration of the malignant leucocytes, autologous cultured dendritic cells are added to the apoptotic T-cells, and the cell mix is incubated overnight with shaking to maximize contact between the T-cells and the dendritic cells. The apoptotic T-cell/dendritic cell mix has proven to be an effective cellular vaccine in test mice challenged with viable tumorigenic 2B4.11 cells.

- 3 -

While the above-described experimental protocol is apparently more efficient and comprehensive than alternative approaches, it requires extensive *ex vivo* cellular manipulations over a period of several days. Accordingly, an *in vivo* procedure which could in a single day provide large numbers of functional dendritic cells and expose those cells to apoptotic tumor cells would greatly simplify the means by which the anti-tumor cellular vaccine could be prepared.

Summary of the Invention

The present invention is based on the convergence of two disparate phenomena: treating monocytes in a manner which induces their differentiation into functional dendritic antigen presenting cells, and treating disease effector agents to render them apoptotic or to inactivate them. By incubating these treated populations together for a period of time sufficient to optimize processing and presentation of disease associated antigens distinctive to the disease effector agents by the dendritic cells, prior to returning the dendritic antigen presenting cells to the patient, clinically enhanced immunity to the disease associated antigens is achieved.

As used herein, the term "disease effector agents" refers to agents that are central to the causation of a disease state in a subject and which express disease-associated antigens. In certain circumstances, these disease effector agents are disease-causing cells which may be circulating in the bloodstream, thereby making them readily accessible to extracorporeal manipulations and treatments. Examples of such disease-causing cells include malignant T-cells, malignant B cells, T-cells and B cells which mediate an autoimmune response, and virally or bacterially infected white blood cells which express on their surface viral or bacterial peptides or proteins. Exemplary disease categories giving rise to disease-causing cells include leukemia, lymphoma, autoimmune disease, graft versus host disease, and tissue rejection. Disease associated antigens which mediate these disease states and which are derived from disease-causing cells include peptides that bind to a MHC Class I site, a MHC Class II site, or to a heat shock protein which is involved in transporting peptides to and from MHC sites (i.e., a chaperone). Disease associated antigens also include viral or bacterial peptides which

- 4 -

are expressed on the surface of infected white blood cells, usually in association with an MHC Class I or Class II molecule.

Other disease-causing cells include those isolated from surgically excised specimens from solid tumors, such as lung, colon, brain, kidney or skin
5 cancers. These cells do not ordinarily circulate in the blood in significant quantity, but can be manipulated extracorporeally in analogous fashion to blood leukocytes, after they are brought into suspension or propagated in tissue culture.

In addition to disease-causing cells, disease effector agents falling within the scope of the invention further include microbes such as bacteria, fungi and viruses
10 which express disease-associated antigens. It should be understood that viruses can be engineered to be "incomplete", i.e., produce distinguishing disease-causing antigens without being able to function as an actual infectious agent, and that such "incomplete" viruses fall within the meaning of the term "disease effector agents" as used herein.

Accordingly, the present invention provides, in one aspect, a method for
15 inducing the differentiation of monocytes contained in an extracorporeal quantity of a subject's blood into functional dendritic antigen presenting cells. According to the invention, the monocytes are treated by at least one of the following: (1) exposing the monocytes to physical perturbation, (2) irradiating the monocytes in the presence of a photoactivatable agent capable of forming photoadducts with cellular components, and
20 (3) treating the monocytes with a DNA binding agent. Following treatment, the monocytes are incubated for a period of time sufficient to maximize the number of functional dendritic cells.

In the preferred embodiment of this aspect of the invention, the monocytes are separated from the extracorporeal quantity of the subject's blood by
25 leukapheresis. Following separation, the monocytes are subjected to photopheresis using 8-MOP and exposure to UV energy. The cells are then incubated for a period of from about 6 to about 48 hours, and most preferably from about 12 to about 24 hours.

In a second aspect, the invention provides a composition comprising functional dendritic antigen presenting cells derived from monocytes treated and

- 5 -

incubated in the manner set forth above. In a preferred embodiment of this aspect of the invention, the monocytes have been separated by leukapheresis and then subjected to photopheresis using 8-MOP and exposure to UV energy. The treated monocytes are incubated for a period of time sufficient to maximize the number of functional dendritic cells in the composition, preferably for a period of from about 6 to 48 hours. The composition preferably further comprises at least one antigen selected for presentation by the dendritic cells and may contain at least one immunomodulatory agent.

The composition may also be provided in packaged form, wherein the composition is stored in a container which does not leach substantial amounts of plasticizer and which is sufficiently porous to permit the exchange of gases, particularly CO₂ and O₂.

The invention provides, in a third aspect, a method of enhancing the presentation of disease associated antigens. According to the method, disease effector agents contained in a quantity of a subject's blood or derived from an exogenous source and capable of expressing at least one disease associated antigen are treated to render the agents apoptotic or to inactivate them. Monocytes contained in a quantity of the subject's blood are also treated by at least one of the following: (1) exposing the monocytes to physical perturbation, (2) irradiating the monocytes in the presence of a photoactivatable agent capable of forming photoadducts with cellular components, and (3) treating the monocytes with a DNA cross-linking agent. The treated disease effector agents and the treated monocytes are then incubated together for a period of time sufficient to induce differentiation of the monocytes into functional dendritic antigen presenting cells and to optimize processing and presentation of the disease associated antigens by the dendritic cells.

In a preferred embodiment of this aspect of the invention, the monocytes and the disease effector agents are separated from the quantity of the patient's blood by leukapheresis. The monocytes and disease effector agents are then exposed to photo-activated 8-MOP and incubated together for a period of from about 6 to about 48 hours,

- 6 -

and preferably for about 12 to about 24 hours. The mix is then reinfused into the subject to elicit an immune response.

In yet another aspect, the invention provides a composition of co-incubated populations comprising a first population including treated disease effector agents which express at least one disease associated antigen, and a second population including functional dendritic antigen presenting cells. The dendritic cells of the second cell population are derived from monocytes which have been treated by at least one of: (1) exposure to physical perturbation, (2) irradiation in the presence of a photoactivatable agent capable of forming photoadducts with cellular components, and (3) treatment with a DNA binding agent.

Preferably, both the monocytes and the disease effector agents are separated from the quantity of the patient's blood by leukapheresis, and both populations are exposed to photo-activated 8-MOP. The first and second populations comprising the composition are co-incubated for a period of from about hours 6 to 40 hours, and the composition preferably includes at least one immunomodulatory agent.

A packaged composition of co-incubated populations is also provided. The packaged composition includes the treated monocytes and treated disease effector agents co-incubated in a container which does not leach substantial amounts of plasticizer and which is sufficiently porous to permit the exchange of gases.

20

Brief Description of the Drawings

Fig. 1 is a graph which illustrates the generation of both dendritic antigen presenting cells and apoptotic T-cells following overnight incubation of blood exposed to 8-MOP and ultraviolet A energy.

Detailed Description of the Invention

25 1. **Differentiation of Monocytes**

As noted above, monocyte differentiation is initiated by exposing monocytes contained in an extracorporeal quantity of a subject's blood to physical perturbation, irradiation in the presence of a photoactivatable agent capable of forming

- 7 -

photoadducts with cellular components, and/or treatment with a DNA binding agent. In the preferred embodiment of the invention, the monocytes are obtained by preparing a white blood cell concentrate in accordance with standard leukapheresis practice using a leukapheresis/photopheresis apparatus of the type well known to those skilled in the art.

5 In addition to monocytes, the white blood cell concentrate also includes lymphocytes and some red blood cells and platelets. Typically, up to two billion white blood cells are collected during leukapheresis. Assuming that monocytes comprise from about 2% to about 50% of the total white blood cell population collected, approximately 40 million to 1 billion monocytes are present in the white blood cell concentrate.

10 Following separation, the monocytes are subjected to photopheresis by exposure to a photoactivatable agent which is capable of forming photoadducts with cellular components, and then irradiating the exposed cells with radiation suitable for activating the agent, typically ultraviolet or visible light. Prior to photopheresis, saline is added to the white blood concentrate to dilute the red blood cell concentration to
15 about 2% by volume, thereby permitting more effective penetration of the activating radiation to the target leukocytes. The photoactivatable agent can be administered to the subject prior to obtaining a quantity of blood from the subject for leukapheresis and photopheresis. Alternatively, or additionally, the photoactivatable agent can be added directly to the extracorporeal bloodstream, typically by injecting the agent into the
20 tubing leading to the leukapheresis/photopheresis apparatus. Regardless of when and how a particular agent is administered, the monocytes must be exposed to the photoactivatable agent for a period of time sufficient for the agent to enter the monocytes and react with cellular components.

Exemplary photoactivatable agents are psoralens, porphyrins, pyrenes,
25 phthalocyanine, retinoid derivatives, photoactivated cortisone, photoactivated antibodies specifically reactive with the monocytes, photactivatable dyes, and monoclonal antibodies which have been linked to porphyrin molecules.

The psoralens are a preferred class of photoactivatable agents for use in the photopheresis procedure. Psoralens are readily absorbed from the digestive track,

- 8 -

reaching peak levels in the blood and other tissues in one to four hours following oral administration, and these agents are excreted almost entirely within 24 hours. Accordingly, the psoralens are particularly suitable for oral administration prior to obtaining an extracorporeal quantity of the subject's blood. The psoralen molecules are inert prior to exposure to irradiation and are transiently activated to an excited state following irradiation. The transiently activated psoralen molecules are capable of forming photoadducts with cellular DNA, proteins or lipids and generating other reactive species, such as singlet oxygen, which are capable of modifying other cellular components, e.g., the cell membrane and cytoplasmic components such as proteins and aromatic amino acids.

The preferred psoralens include 8-methoxypsoralen (8-MOP), 4' aminomethyl-4, 5', 8 trimethyl-psoralen (AMT), 5-methoxypsoralen (5-MOP), and trimethyl-psoralen (TMP). 8-MOP is the most preferred photoactivatable agent for use with the methods of the invention, and the conditions for oral administration of this psoralen are described in U.S. Patent No. 5,147,289, the disclosure of which is incorporated herein by reference.

The irradiation stage of photopheresis is performed by passing the monocyte/lymphocyte fraction through an exposure device which may be contained within the leukapheresis/photopheresis apparatus or may be physically separate. The preferred exposure device includes a transparent plastic channel having a diameter of about 1 mm disposed between opposed irradiation sources. However, it may also be a compressible transparent compartment, which can be shaken to evenly expose the suspended cells to the light. In this manner, under certain circumstances, cellular adhesion to plastic or excessive disruption of certain types of disease effector cells can be minimized. Referring again to the preferred embodiment, as the monocyte/lymphocyte fraction passes through the channel, the cells are never separated from the irradiation sources by more than about 0.5 mm of blood. Maintaining the monocytes in such close proximity to the irradiation sources has proven particularly effective in ensuring adequate exposure of the monocyte/lymphocyte fraction to the

- 9 -

activating radiation. In the case where a psoralen such as 8-MOP is used as the photoactivatable agent, the irradiation sources emit ultraviolet A radiation (UVA) as the activating radiation. To activate the psoralen, the monocytes are typically exposed to about 1-2 joules/cm² of UVA for a period of from about 15 to about 150 minutes.

5 As noted above, other methods of initiating the differentiation of monocytes to functional dendritic cells may be employed. It has been found that differentiation can be initiated simply by exposing the monocytes to physical perturbation. Thus, for example, centrifuging the monocytes during leukapheresis, and/or passing the monocytes through the narrow channel in the exposure device,
10 thereby applying gentle shearing forces to them, or subjecting the monocytes to hydrostatic pressure will initiate differentiation of the monocytes into functional dendritic cells. This differentiation into dendritic antigen presenting cells is maximized after about 6 to about 48 hours incubation, under conditions described in this application. Differentiation can also be initiated by administering a DNA binding
15 agent, such as mitomycin C or a cis-platinum compound, to the subject. While these agents may be effective in initiating differentiation, they remain in an active state when returned to the subject and thus are not as desirable as the psoralens for treating the monocytes.

 It should also be understood that it is not absolutely necessary to
20 separate the monocytes from the extracorporeal quantity of the patient's blood prior to treatment. Subjecting the extracorporeal quantity of the patient's blood to physical forces, or administering the photoactivatable agent to the subject prior to photopheresis and/or adding the agent to the quantity of blood, and then exposing the quantity of blood including the monocytes to the activating radiation are also effective in initiating
25 differentiation of the monocytes into functional dendritic cells. As long as the monocytes are sufficiently exposed to physical forces, and/or pharmacologic agents, such as photo-activated 8-MOP, to initiate differentiation into dendritic cells followed by subsequent incubation, separation of the monocyte population is not required.

- 10 -

Following treatment to initiate differentiation, the treated monocytes are sequestered for incubation. A standard blood bag may be utilized for this purpose, as is typical in photopheresis. However, it has been found to be particularly advantageous to use a blood bag of the type which does not leach substantial amounts of plasticizer and which is sufficiently porous to permit exchange of gases, particularly CO₂ and O₂. Such bags are available from, for example, the Fenwall division of Baxter Healthcare Corp. under the name Amicus™ Apheresis Kit. Various plasticizer-free blood bags are also disclosed in U.S. Patent Nos. 5,686,768 and 5,167,657, the disclosures of which are herein incorporated by reference.

The treated monocytes are incubated for a period of time sufficient to maximize the number of functional dendritic cells in the incubated cell population. Typically, the treated monocytes are incubated for a period of from about 6 to about 48 hours, with the preferred incubation time extending over a period of from about 12 to about 24 hours. By treating monocytes in the manner described above and then incubating the treated cell population for a minimum of from about 6 to to about 12 hours, and preferably from about 12 to about 40 hours, large numbers of functional dendritic cells are obtained. It has been found to be particularly advantageous to add a buffered culture medium to the blood bag and one or more cytokines, such as GM-CSF and IL-4, during the incubation period.

The mechanism by which the methods of the present invention induce monocyte differentiation into functional dendritic cells is presently under investigation. One possibility is that cytokines liberated from the treated mononuclear cells, including GM-CSF and IL-4, cause the maturation step. In the case where the monocytes are treated with 8-MOP, it may be that 8-MOP binding increases the cytoplasmic cAMP level, which is known to enhance monocyte-dendritic cell differentiation. It is also possible that other factors, either individually or collectively, have this impact, or that 8-MOP directly or indirectly through its DNA and/or protein binding has this effect. Whatever the underlying mechanism, inducing monocyte differentiation according to the invention provides dendritic cells in numbers which equal or exceed the numbers of

- 11 -

dendritic cells that are obtained by expensive and laborious culture of leukocytes in the presence of cytokines such as GM-CSF and IL-4 for seven or more days.

The large numbers of functional dendritic cells generated during incubation provide a ready means of presenting selected disease associated antigens and are thereby conducive to efficient immunotherapy. Antigen preparations selected to elicit a particular immune response and derived from, for example, tumors, disease-causing non-malignant cells, or microbes such as bacteria, viruses and fungi, can be added directly to the blood bag during incubation. The microbes may preferably be inactivated by prior exposure to 8-MOP or other agents. It is known that 8-MOP can cause apoptosis in bacteria and fungi and can inactivate viruses. Bringing mature dendritic cells into close contact with such antigen preparations within the confines of the blood bag provides large numbers of antigen-loaded dendritic cells. The antigen-loaded dendritic cells can be used as immunogens by reinfusing the cells into the subject or by otherwise administering the cells in accordance with methods known to elicit an immune response, such as subcutaneous, intradermal or intramuscular injection. As described below, it is also possible to generate antigen-loaded dendritic cells by treating and co-incubating monocytes and disease effector agents which are capable of expressing disease associated antigens.

2. Methods for Enhancing the Presentation of Disease Associated Antigens.

This aspect of the invention relates in general to improved immunotherapeutic methods for treating disease states mediated by disease effector agents. As discussed above, such agents comprise microbes, such as bacteria, fungi, and complete and incomplete viruses, and disease-causing clonal populations of cells, including clones of malignant cells or clones of non-malignant T- or B-cells attacking the individual's own tissues or transplanted tissues. Since these agents have distinctive antigens on their surface that permit them to be distinguished from most other cells, immune reactions can be ideally developed against their distinctive antigens. These immune reactions can then suppress or eliminate the disease effector agent populations. Through the generation of dendritic antigen-presenting cells capable of effectively

- 12 -

introducing the relevant antigens to a responding immune system, this invention substantially enhances the likelihood of such a disease-controlling immunologic response.

Central to this aspect of the invention is the co-cultivation of increased
5 numbers of dendritic antigen presenting cells, generated as described above, with clones of apoptotic disease-causing cells or inactivated or incomplete microbes which bear distinctive antigens. In the case of disease-causing cells, bacteria and fungi, other means of inducing apoptosis, in addition to exposure to photo-activated drugs, may be applicable.

10 For example, synthetic peptides with the arginine-glycine-aspartate (RGD) motif could be added to cell suspensions of the disease-causing cells isolated from the patient's blood, from excised solid tumors or tissue cultures of the same. RGD has been shown (Nature, Volume 397, pages 534-539, 1999) to induce apoptosis in tumor cells, possibly by triggering pro-caspase-3 autoprocessing and activation.
15 Similarly, apoptosis could be induced in cells having Fas receptors, by stimulating with antibodies directed against this receptor, in this way sending signals to the inside of the cell to initiate programmed cell death, in the same way that normally Fas ligand does. In addition, apoptosis can be induced by subjecting disease-causing cells to heat or cold shock, certain viral infections (i.e., influenza virus), bacterial toxins, and x-ray or
20 gamma-irradiation. Alternatively, certain infectious agents such as influenza virus can cause apoptosis and could be used to accomplish this purpose in cell suspensions of disease-causing cells.

Hence, these approaches, although not as usually preferred as the induction of apoptosis by photo-activated 8-MOP, could accomplish the purpose of
25 initiating apoptosis or inactivation in disease-causing cellular populations, prior to their co-cultivation with the induced dendritic antigen-presenting cells and return to the patient for purposes of immunization. Of course, it should be understood that since viruses are not cells, they cannot undergo apoptosis as that term is generally understood and used by those skilled in the art. It is known, however, that viruses can be

- 13 -

inactivated by exposure to 8-MOP and other photo-activated drugs and therefore can be treated in this manner prior to their co-cultivation with induced dendritic antigen presenting cells.

This aspect of the invention will be described with particular reference
5 to an enhanced therapy for treating cutaneous T-cell lymphoma. However, it should be understood that the invention is not limited to this particular application and that the invention may be employed to treat any disease state which includes as a component disease effector agents distinguishable by their own surface antigens. A number of such disease states, component effector agents and disease associated antigens have
10 been discussed above.

Cutaneous T-cell lymphoma (CTCL) is an immune disease that is caused by a massive expansion of a single clone of aberrant T-cells. These malignant cells are distinguished by clone-specific or tumor-specific cell surface antigens, at least one set of which are derived from clone specific protein components of the clone-specific T-cell
15 receptor. Cytotoxic T-cell responses can be generated selectively against these clone-specific antigens. During the past decade, photopheresis has become a standard immunotherapy for advanced CTCL and works, at least in part, by generating such anti-CTCL immune responses. In standard CTCL treatment using photopheresis, leukocytes and monocytes are separated by leukapheresis from an extracorporeal
20 quantity of a subject's blood. The monocytes and leukocytes are circulated through an ultraviolet A exposure system of the type described above, in which biologically inert 8-MOP is activated to covalently bond to DNA and cytoplasmic proteins in the lymphocytes and monocytes. This is a highly directed therapy, since the drug remains active for only millionths of a second, thereby chemically altering only those cells in
25 the exposure field and explaining the paucity of systemic side effects. Photopheresis provides increased immunogenicity of the exposed leukocytes, without causing general immunosuppression. Thus, returning the treated cells to the subject can lead to a "vaccination" effect which, in the most responsive subjects, results in a sustained immunologic response to the chemically altered and reintroduced leukocytes.

- 14 -

Alteration and return of less than 5% of the body burden of malignant T-cells can induce a meaningful anti-tumor response which in some subjects has resulted in complete remissions lasting more than fifteen years. Methods for applying photopheresis to the treatment of CTCL are disclosed in U.S. Patent Nos. 5,114,721 and 4838,852 and published PCT applications WO 97/34472 and WO 94/11016, the disclosures of which are incorporated herein by reference.

The clinical results achieved through the application of photopheresis to CTCL have encouraged a search for the treatment's underlying mechanism for two major reasons. First, if the mechanism by which photopheresis vaccinates patients against their malignant cells could be better understood, it should then be possible to refine the methodology and enhance its efficacy. For example, only 25% of the patients with advanced CTCL have a major persistent response to photopheresis. While these positive responses are profound and their frequency exceeds those produced by prior conventional chemotherapy, it would be desirable to increase the efficiency of the procedure. Second, if the mechanism could be better understood, it should then also be possible to extend the revised therapy to other types of malignancies and disease processes. This application is based on the new recognition of the role of dendritic antigen presenting cells in the response to photopheresis, and more particularly on methodology of enhancing this role. Studies in experimental systems and with transformed human cells lines have yielded four lines of evidence. First, the treatment stimulates CD8 T-cells to suppress the activity of pathogenic clones of T-cells. Second, these CD8 cells, at least in CTCL where there is only a single clone of pathogenic T-cells, recognize tumor-specific peptide antigens in the context of Class I MHC complexes at the tumor cell surface. Third, exposure of human lymphoblasts to photo-activated 8-MOP triples the display of Class I complexes, peaking after overnight incubation. Finally, the treatment also causes apoptosis in lymphocytes and their ingestion by phagocytic mononuclear cells.

Multiple lines of clinical and experimental evidence have confirmed the "vaccination" phenomenon which is associated with the induction of potent CD8

- 15 -

responses capable of selectively suppressing aberrant T-cell populations. In the case of CTCL, at least some of the anti-cancer CD8 T-cells selectively targeted tumor-specific peptides derived from the T-cell receptor proteins of the malignant cells. Since the T-cell receptors of CD8 T-cells recognize antigenic peptides in the context of Class I MHC, attention has focused on the impact of 8-MOP on the display of these complexes. It has recently been reported that 8-MOP triples the display of Class I at the cell surface of transformed human lymphocytes, maximizing about 22 hours after exposure, and that this effect is dependent on the degradation of cytoplasmic proteins and the transport of the generated peptide fragments across the endoplasmic reticulum through TAP pores. This effect appears to be initiated by binding of 8-MOP to aromatic amino acids of cytosolic proteins rather than the drug's other main molecular target, pyrimidine bases of DNA.

The present invention is based on the assumption that if an immune response is to be generated against weakly immunogenic complexes containing the relevant antigens, then such a response might be maximized if the complexes are maximized on the antigen presenting cells. In conventional photopheresis, T-cells are immediately returned to the subject at a point when apoptosis is only modestly elevated over baseline and when Class I complexes are also only modestly enhanced. In the present method, the treated leukocytes are incubated overnight, typically for a period of from about 6 to 48 hours. An unexpected finding was that overnight incubation of the treated cells not only enhances the expression of Class I complexes by the apoptotic T-cells, but also maximized the maturation of monocytes into functional dendritic cells. Thus, the convergence of these two phenomena made the incubation phase a simple means of bringing large numbers of apoptotic malignant cells into apposition with increased numbers of functional dendritic cells capable of ingesting apoptotic cells or fragments of apoptotic cells. It has previously been shown the mononuclear cells in the photopheresis bag have already begun to phagocytose apoptotic T-cells, although these mononuclear cells do not have the properties of dendritic cells. Typically, antigen presenting cells process endocytosed antigens through the Class II MHC pathway,

- 16 -

which ordinarily stimulates expression of CD4T-cells rather than the desired CD8 cytotoxic cells which "see" antigens only in the context of Class I MHC. However, it is important to note that it has recently been reported that dendritic cells have a special capacity to process and present antigens derived from apoptotic cells through the Class
5 I MHC system.

An enhanced photopheresis protocol based on the present invention has provided encouraging clinical results in a pilot study which included four subjects suffering from advanced CTCL. However, before discussing the clinical results of the study, a treatment protocol embodying the present invention will be set forth in the
10 following examples.

Example I: Photopheresis Protocol

The first step, which is the photopheresis protocol, is essentially the same as the protocols currently approved by the FDA. Subjects receive either oral 8-MOP (0.6 mg/kg) or intravenous 8-MOP directly into the photopheresis apparatus, to
15 yield a concentration of 50-200 ng/ml of drug. Next, the blood is leukapheresed to obtain a buffy coat and is then passed through a contiguous closed circuit ultraviolet A exposure device, which delivers about 1-2 joules /cm² of ultraviolet A energy (320 nm-400 nm). In this manner, about 1 to 100 molecules of 8-MOP are induced to covalently bind to each million base pairs of DNA. A nearly equal amount of 8-MOP is induced
20 to covalently bind to aromatic amino acids of cytoplasmic proteins. The treated leukocyte fraction, comprising a total volume of approximately 250 cc, is combined with 500 cc saline and then sequestered in a standard blood bank bag, as is typical for the photopheresis procedure. Following photopheresis, the treated fraction is subjected to the following novel incubation phase protocol.

Example II: Incubation Phase Protocol

Following collection of the post photopheresis sample after ultraviolet A activation with 8-MOP, the treated cell populations are incubated as follows:

- 17 -

1. Remove two Amicus platelet storage bags (Baxter Fenwall PL 2410) from an apheresis kit (Baxter Fenwall 4R 23-12) by heat sealing the tubing and cutting the tubing at the end connecting to the kit.
2. Insert a sharp catheter into the pheresis bag (spike), thereby
5 breaking the seal, with a Charter Medical 3-leg transfer set (#O3-220-02) and clamp the tubing. Spike the two Amicus bags with the other piercing pins of the same transfer set thereby establishing a passageway for the transfer of the cell suspension.
3. Hang the pheresis bag on an IV pole and open the clamp
10 allowing 1/2 of the pheresis to drain into each Amicus bag by gravity, and then clamp the tubing.
4. Remove the spikes and replace with sampling site couplers.
5. Place each Amicus bag in a separate Fenwall centrifuge bag and
into a centrifuge carrier.
6. Centrifuge for 10 minutes, at 1000 rpm, 23_C, to concentrate the
15 cells as a pellet at the bottom of each bag to permit removal of a large fraction of the plasma, which contains traces of plasticizer.
7. After centrifugation, insert a needle attached to the tubing on a transfer pak into the sampling coupler on one of the Amicus bags.
8. Carefully place the Amicus bag in a plasma extractor to avoid
20 resuspending the cell pellet. Close the extractor and express the plasma into the transfer bag by slowly tipping the extractor forward. When approximately 50 cc has drained into the transfer bag and/or the pellet begins to resuspend, return the extractor to an upright position and remove the needle.
9. Remix the contents of the bags by gentle agitation being careful
25 to resuspend any adherent cells attached to the bag wall.
10. Spike one 500 cc bottle containing 100 cc of colorless RPMI 1640 media with Hepes Buffer with a Baxter vented medication set and clamp the tubing. Insert the attached needle into the sampling coupler port on the first Amicus

- 18 -

bag. Hang the bottle on the IV pole and open the tubing allowing the media to drain into the bag.

11. Clamp the tubing and remove the needle and discard the medication set. Mix the bag by gentle inversion and place the bag in a 370_C incubator
5 on a shelf with the Abel side down, overnight.

12. Repeat steps 8-12 for the second bag.

13. Following incubation for a period of __ hours, remove one bag from the incubator, gently mix by agitation and inversion, making sure that all adherent cells are resuspended. Take out 60 cc of blood in a syringe. Inject one
10 aerobic, and one anaerobic blood culture bottle for microbiology. Inject one lavender top tube for WBC and differential to be sent to hematology.

14. Resuspend the second Amicus bag and place both bags in individual centrifuge bags and centrifuge.

15. Remove and transfer the supernatant fluid as described in steps
15 8-10.

16. Return well mixed blood to the patient.

Fig. 1 is a composite graph which illustrates the generation of both dendritic antigen presenting cells and apoptoticT-cells following treatment by the photopheresis and incubation protocols set forth above. As shown in Fig. 1, pre-
20 treatment blood contained nearly undetectable numbers of dendritic cells, using either the $\alpha V\beta 5$ or CD11c markers for identification. After incubation for about 22 hours, both of these markers revealed large numbers of mature dendritic cells. Similarly, the pre-treated blood contained very few apoptoticT-cells. Only after overnight incubation
25 did apoptoticT-cells become significantly evident, as illustrated by the simultaneous identification of theT-cells with the CD3 marker and the apoptotic cells with the APO2 markers.

The fourth set of bars at the far right of the graph illustrates the differentiation of monocyte into mature dendritic cell by means of physical perturbation

- 19 -

and incubation only, without exposure to ultraviolet light. Differentiation was initiated by isolating monocytes and T-cells from an extracorporeal quantity of blood by leukapheresis. The isolated monocytes and T-cells were not subjected to photopheresis but were exposed only to the centrifugal forces associated with leukapheresis. The isolated cell populations were then incubated for a period of about 22 hours according to the incubation protocol set forth above. As shown in Fig. 1, the physical forces applied during leukapheresis, together with overnight incubation, caused the monocytes to efficiently evolve into functional dendritic cells, as identified by the α V β 5 and CD11c markers. No significant apoptosis of the T-cells was observed, indicating that treatment with 8-MOP followed by exposure to UV, or some other form of treatment as described above, is required to induce T-cell apoptosis.

The Y axis of the graph gives the number of functional dendritic cells per cubic centimeter. Since the total volume incubated over the 22 hour period was 250 cc, 32.5 million dendritic cells (130,000 x 250) were generated, as indicated in the third set of bars by the CD11c marker. It has been shown that dendritic cells having this level of maturity phagocytose apoptotic cells and are efficient presentors of antigens derived from such cells. Monocytes may also ingest apoptotic cells or fragments of such cells, but monocytes cannot efficiently present antigen material processed from the apoptotic cells to CD8 cytotoxic T cells. CD8 T cells only recognize antigens which are associated with Class I MHC at the surface of the antigen presenting cell. Monocytes primarily present antigens derived from ingested cells in association with Class II MHC molecules, which CD8 T cells cannot recognize. Dendritic cells, on the other hand, in part because they include the α V β 5 integrin, have the special ability to "cross-prime" CD8 T cells by presenting the antigens derived from the digestion of apoptotic cells and displaying the processed antigens in association with the Class I MHC molecules that CD8 cytotoxic T cells can recognize. This is a major reason why functional dendritic cells are so useful in stimulating tumor immunity, or suppressing undesirable immunologic processes by attacking the aberrant T cells that cause them.

- 20 -

The graph illustrated in Fig. 1 further demonstrates an effective means of determining the optimum incubation time for the mixed cell populations. Since the particular markers employed permit the numbers of dendritic cells and apoptotic T-cells to be quantified simultaneously, the incubation time that results in the optimal combination of apoptotic cells and newly formed dendritic cells can be readily determined. This is the controlling determinant establishing when to terminate incubation and reinfuse the incubated cells into the subject.

As noted previously, an incubation time of from about 12 to about 48 hours results in a maximum number of dendritic cells. The apoptotic T-cells maximize in a period of about 6 to about 40 hours. Accordingly, an incubation period of from about 6 to about 24 hours provides the most advantageous combination of apoptotic T-cells and dendritic cells. After an incubation period of this duration, the number of apoptotic cells is at a maximum and large numbers of functional dendritic cells are also present in the incubation bag. Thus, a maximum number of apoptotic cells capable of expressing disease-associated antigens are present and a large number of functional dendritic cells capable of processing and presenting those antigens are also present. In the case where the disease effector agent is derived from an exogenous source and is added to the incubation bag, the incubation period required for maximizing the number of apoptotic cells is obviously not a factor. In such instances, the time period required for maximizing the number of induced dendritic cells is the factor which determines the duration of the co-incubation.

3. Clinical Efficacy of Combined Treatment and Co-Incubation

The treatment method taught by the present invention has been tested in a pilot study involving four CTCL subjects whose disease had been advancing while on standard photopheresis. The four patients in the pilot study were carefully selected from a large CTCL population based on three criteria: (1) increasing tumor burden despite continued conventional photopheresis; (2) malignant clones that could be quantified in blood; and (3) low absolute blood CD8 levels. The leukemic cells in three of the subjects could readily be distinguished from normal T-cells, since their clonal

- 21 -

T-cell receptor phenotype was recognizable using fluorescein-tagged anti-family V T-cell receptor monoclonal antibodies (V mAb). Values above 5% indicate expansion of the malignant clone. Although the clonal T-cell receptor of the fourth patient's CTCL cells does not bind any currently available V mAb, the CD4/CD8 ratio permits
5 quantitation of that patient's leukemic population as well. The unresponsiveness of the four patients to conventional photopheresis likely reflects their CD8 T cell deficiency, since clinical responders usually require an intact CD8 compartment. Therefore, these patients present a significant challenge for the new treatment approach. Although the study population was small, it was easy to quantify reversal of disease progression in
10 this poor prognosis patient group.

Following treatment with the above-described protocol, each of the four patients had a diminution in the absolute circulating malignant pool over the twelve months of the protocol. Whereas none experienced complete hematologic remission, the previous rapid increases in blood CTCL cells were reversed. Those
15 symptomatic infections common in individuals whose immune systems have been compromised by their CTCL, and the therapy for this disease, were not encountered. Measurements of tumor burden and clinical response centered on blood determinations and quantitation of the number of infiltrating T cells in biopsies of the clinically most severe skin lesions. It is important to note that the severity and distribution of skin
20 lesions in three of the four patients lessened. In one patient, long-standing, maximal, generalized exfoliative erythroderma associated with intractable pruritus was transformed to low grade, nearly asymptomatic erythroderma, and two of the other patients had nearly complete cutaneous remissions.

The photopheresis/incubation protocol tested in this study, like
25 conventional photopheresis, appears to be safe, since no side effects were encountered in these subjects. Further, the capacity of the protocol to bring together malignant apoptotic cells bearing the relevant immunizing antigens with functional dendritic cells capable of presenting these antigens to a responding immune system offers additional opportunities for immunotherapy beyond the treatment of CTCL. For example, in a

- 22 -

recently reported randomized, controlled trial, the combination of photopheresis with conventional immunosuppressive drugs proved effective in reducing the number of rejection episodes experienced by heart transplant recipients. Preliminary studies have also suggested the efficacy of conventional photopheresis in certain autoimmune diseases, such as rheumatoid and psoriatic arthritis, lupus erythematosus, scleroderma and graft-versus-host disease (following allogeneic bone marrow transplantation). The present invention's capacity to provide an *in vivo* source of large numbers of dendritic cells should enhance these therapies. Modifications to the protocol may also permit co-cultivation of dendritic cells derived from induced monocytes with suspended apoptotic solid tumor cells, apoptotic infectious microbes or inactivated or incomplete viruses.

Accordingly, it should be understood, as noted above, that while certain aspects of the invention have been described in connection with an enhanced therapy for CTCL, the invention is applicable to a broad range of immune diseases without departing from the spirit and scope of the invention.

- 23 -

I CLAIM:

1. A method for inducing differentiation of monocytes contained in an extracorporeal quantity of a subject's blood into functional dendritic antigen presenting cells, said method comprising the steps of:

(a) treating the monocytes by at least one of the following: (1) exposing the monocytes to physical perturbation, (2) irradiating the monocytes in the presence of a photoactivatable agent capable of forming photoadducts with cellular DNA components, and (3) treating the monocytes with a DNA binding agent; and

(b) incubating the treated monocytes for a period of time sufficient to maximize the number of functional dendritic cells.

2. The method of claim 1, wherein prior to step (a) the method further comprises the step of:

separating the monocytes from the extracorporeal quantity of the subject's blood by subjecting the quantity of blood to a leukapheresis process.

3. The method of claim 2, wherein the monocytes are incubated for a period of from about 6 to about 48 hours.

4. The method of claim 3, wherein the monocytes are incubated for a period of from about 12 to about 24 hours.

5. The method of claim 3, wherein the monocytes are irradiated in the presence of the photoactivatable agent.

6. The method of claim 6, wherein the photoactivatable agent is 8-MOP.

7. The method of claim 3, wherein the monocytes are exposed to the DNA binding agent.

8. The method of claim 3, wherein the treated monocytes are incubated together with at least one of GM-CSF and IL-4.

9. The method of claim 3, wherein the monocytes are incubated in a container which does not leach substantial amounts of plasticizer and which is sufficiently porous to permit exchange of gases.

- 24 -

10. The method of claim 3, wherein the treated monocytes are incubated together with at least one selected antigen to be processed and presented by the dendritic cells.

11. The method of claim 10, wherein the at least one antigen is expressed on the surface of a disease effector agent.

12. The method of claim 11, wherein the disease effector agent is selected from the group consisting of disease-causing cells and microbes.

13. A composition comprising functional dendritic antigen presenting cells derived from monocytes which have been incubated following their treatment by at least one of: (1) exposure to physical perturbation, (2) irradiation in the presence of a photoactivatable agent capable of forming photoadducts with cellular components, and (3) treatment with a DNA binding agent.

14. The composition of claim 13, wherein the treated monocytes have been incubated for a period of time sufficient to maximize the number of functional dendritic cells present in the composition.

15. The composition of claim 14, wherein the treated monocytes have been incubated for a period of from about 6 to about 48 hours.

16. The composition of claim 15, wherein the monocytes have been incubated for a period of from about 12 to about 24 hours.

17. The composition of claim 14, wherein the monocytes have been irradiated in the presence of the photoactivatable agent.

18. The composition of claim 17, wherein the photoactivatable agent is 8-MOP.

19. The composition of claim 14, wherein the monocytes have been exposed to the DNA binding agent.

20. The composition of claim 14, further comprising at least one of GM-CSF and IL-4.

21. The composition of claim 14 further comprising at least one selected antigen for presentation by the dendritic cells.

- 25 -

22. The composition of claim 21, wherein the at least one antigen is expressed on the surface of a disease effector agent.

23. The method of claim 22, wherein the disease effector agent is selected from the group consisting of disease-causing cells and microbes.

24. A packaged preparation comprising:

a composition including functional dendritic antigen presenting cells derived from monocytes which have been incubated following their treatment by at least one of exposure to physical perturbation, irradiation in the presence of a photoactivatable agent capable of forming photoadducts with cellular components, and treatment with a DNA binding agent; and

a container which does not leach substantial amounts of plasticizer and which is sufficiently porous to permit exchange of gases for storing the composition.

25. The packaged preparation of claim 24, wherein the treated monocytes have been incubated for a period of time sufficient to maximize the number of functional dendritic cells present in the composition.

26. The packaged preparation of claim 25, wherein the composition further includes at least one of GM-CSF and IL-4.

27. The packaged preparation of claim 25, wherein the composition further includes at least one selected antigen for presentation by the dendritic cells.

28. A method of enhancing the presentation of disease associated antigens, said method comprising the steps of:

(a) treating disease effector agents capable of expressing at least one disease associated antigen to one of: render the agents apoptotic and inactivate the agents;

(b) treating monocytes contained in an extracorporeal quantity of a subject's blood by at least one of the following: (1) exposing the monocytes to physical perturbation, (2) irradiating the monocytes in the presence of a photoactivatable agent capable of forming photoadducts with cellular components, and (3) treating the monocytes with a DNA binding agent; and

- 26 -

(c) incubating the treated disease effector agents and the treated monocytes together for a period of time sufficient to induce differentiation of the monocytes into functional dendritic antigen presenting cells and to optimize processing and presentation of the disease associated antigens by the dendritic cells.

15 29. The method of claim 28, wherein the treated disease effector agents and the treated monocytes are incubated together for a period of from about 6 to about 40 hours.

 30. The method of claim 29, wherein the treated disease effector agents and the monocytes are incubated together for a period of from about 12 to about
20 24 hours.

 31. The method of claim 28, wherein the disease effector agents are contained in the extracorporeal quantity of the subject's blood.

 32. The method of claim 28, wherein the disease effector agents are derived from an exogenous source.

25 33. The method of claim 31, wherein prior to step (a) the method further comprises the step of:

 separating the disease effector agents and the monocytes from the extracorporeal quantity of the subject's blood by subjecting the quantity of blood to a leukapheresis process.

 34. The method of claim 28, wherein the disease effector agents are selected from the group consisting of disease-causing cells and microbes.

 35. The method of claim 34 wherein the disease-causing cells selected from the group consisting of T-cells, B-cells and macrophages.

 36. The method of claim 35, wherein the T-cells include lymphoma cells.

 37. The method of claim 36, wherein the lymphoma cells include cutaneous T-cell lymphoma cells.

- 27 -

38. The method of claim 28, wherein steps (a) and (b) are performed simultaneously by irradiating the disease effector agents and the monocytes with a photoactivatable agent capable of forming photoadducts with DNA and cell proteins.

39. The method of claim 38, wherein the photoactivatable agent is 8-MOP.

40. The method of claim 28, wherein the disease associated antigen is selected from the group consisting of viral antigens, fungal antigens, bacterial antigens, transplant antigens and tumor specific antigens.

41. The method of claim 40, wherein the tumor specific antigen is a peptide derived from a T-cell antigen-binding cell surface receptor.

42. The method of claim 28, wherein the disease effector agents and the monocytes are incubated together with at least one of GM-CSF, IL-4, TNF-alpha, and IL-12.

43. The method of claim 28, wherein the disease effector agents and the monocytes are incubated together in a container which does not leach substantial amounts of plasticizer.

44. The method of claim 28, further including the step of:
administering the incubated disease effector agents and the dendritic cells to the subject to elicit an immune response.

45. The method of claim 44, wherein the incubated disease effector agents and the dendritic cells are administered to the subject in at least one of the presence of and in combination with an immunomodulatory agent.

46. A composition of co-incubated populations comprising:
a first population including disease effector agents which express at least one disease associated antigen; and

a second population including functional dendritic antigen presenting
5 cells derived from monocytes which have been treated by at least one of: (1) exposure to physical perturbation, (2) irradiation in the presence of a photoactivatable agent

- 28 -

capable of forming photoadducts with cellular components, and (3) treatment with a DNA binding agent.

47. The composition of claim 46, wherein the first and second populations have been co-incubated for a period of from about 6 to about 40 hours.

48. The composition of claim 47, wherein the first and second populations have been co-incubated for a period of from about 12 to about 24 hours.

49. The composition of claim 46, wherein the disease effector agents are selected from the group consisting of disease-causing cells and microbes.

50. The composition of claim 49, wherein the disease-causing cells are selected from the group consisting of T-cell, B-cells and macrophages.

51. The composition of claim 50, wherein the T-cells include lymphoma cells.

52. The composition of claim 51, wherein the lymphoma cells include cutaneous T-cell lymphoma cells.

53. The composition of claim 46, wherein the disease effector agents and the monocytes have been treated with a photoactivatable agent.

54. The composition of claim 53, wherein the photoactivatable agent is 8-MOP.

55. The composition of claim 46, wherein the disease effector cells and the monocytes have been treated with a DNA binding agent.

56. The composition of claim 46, wherein the monocytes are separated from an extracorporeal quantity of a subject's blood.

57. The composition of claim 56, wherein the disease effector agents are one of contained in the extracorporeal quantity of the subject's blood and derived from an exogenous source.

58. The composition of claim 46, wherein the disease associated antigen is selected from the group consisting of viral antigens, fungal antigens, bacterial antigens, transplant antigens, and tumor specific antigens.

- 29 -

59. The composition of claim 46, further including at least one immunomodulatory agent.

60. A packaged composition of co-incubated populations comprising:

a first population including disease effector agents which express at least one disease associated antigen;

5 a second population including functional dendritic antigen presenting cells derived from monocytes which have been treated by at least one of: (1) exposure to physical perturbation, (2) irradiation in the presence of a photoactivatable agent capable of forming photoadducts with cellular components, and (3) treatment with a DNA binding agent; and

10 a container which does not leach substantial amounts of plasticizer and which is sufficiently porous to permit exchange of gases for storing the composition.

61. A method of providing a vaccine against disease effector agents capable of expressing at least one disease associated antigen, said method comprising the steps of:

(a) treating the disease effector agents to one of: render the agents apoptotic and inactivate the agents;

(b) treating monocytes contained in an extracorporeal quantity of a subject's blood by at least one of the following: (1) exposing the monocytes to physical perturbation, (2) irradiating the monocytes in the presence of a photoactivatable agent capable of forming photoadducts with cellular components, and (3) treating the monocytes with a DNA binding agent;

(c) incubating the treated disease effector agents and the treated monocytes together for a period of time sufficient to induce differentiation of the monocytes into functional dendritic antigen presenting cells and to optimize processing and presentation of the disease associated antigens by the dendritic cells; and

15 (d) administering the incubated disease effector agents and the dendritic cells to the subject to vaccinate the subject against the disease effector agents.

- 30 -

62. The method of claim 61, wherein the disease effector agents are selected from the group consisting of disease-causing cells and microbes.

63. The method of claim 61, wherein the disease effector agents are one of: contained in the extracorporeal quantity of the subject's blood and derived from an exogenous source.




FIG. 1

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/08793

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	SCHEDING et al. FLT-3 ligand, TGF- β and GM-CSF/IL-4 are Critical Growth Factors for the Induction of CD1a+/CD14-/CD80+ Dendritic Cell (DC) Development from CD34+ PBPC in Serum-Free Medium. Journal of the American Society of Hematology. 15 November 1997, Vol. 90, No. 10, Supplement 1, Abstract # 2125, see abstract.	1-63
Y	STROBL et al. TGF- β 1 Promotes In Vitro Development of Dendritic Cells from CD34+ Hemopoietic Progenitors. J. Immunology. 15 August 1996, Vol. 157, No. 4, pages 1499-1506, see entire article.	1-63
Y	RIEDL et al. TGF- β 1 Promotes In Vitro Generation of Dendritic Cells by Protecting Progenitor Cells from Apoptosis. J. Immunology. 15 February 1997, Vol. 158, No. 4, pages 1591-1597, see entire article.	1-63
Y	THOMAS et al. Dendritic Cells: Origin and Differentiation. Stem Cells. 1996, Vol. 14, pages 196-206, see entire article.	1-63
Y	JAFFE, R. Review of Human Dendritic Cells: Isolation and Culture from Precursors. Pediatric Pathology. 1993, Vol. 13, pages 821-837, see entire article.	1-63
Y	WILLIAMS et al. Isolation and Function of Human Dendritic Cells. International Review of Cytology. 1994, Vol. 153, pages 41-103, see entire article.	1-63

INTERNATIONAL SEARCH REPORT

 International application No.
 PCT/US00/08793

A. CLASSIFICATION F SUBJECT MATTER IPC(7) : A61K 48/00; C12N 5/00, 5/10 US CL : 424/93.1, 93.21; 435/325, 343, 375, 383, 372 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 424/93.1, 93.21; 435/325, 343, 375, 383, 372; 514/44 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) WEST, STN, MEDLINE, EMBASE, BIOSIS, CAPLUS search terms: dendritic cells, GM-CSF, IL-4, monocytes, DNA, binding		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	STROBL et al. flt3 Ligand in Cooperation with Transforming Growth Factor- β 1 Potentiates In Vitro Development of Langerhans-Type Dendritic Cells and Allows Single-Cell Dendritic Cell Cluster Formation Under Serum-Free Conditions. Blood. 15 August 1997, Vol. 90, No. 4, pages 1425-1434, see entire article.	1-63
Y	GARBE et al. Transforming Growth Factor Beta (TGF- β 1) Synergizes in the Differentiation of Dendritic Cells from Monocytes Under Serum-Free Culture Conditions but Inhibits Upregulation of Costimulatory Molecules and MHC Class II. Journal of the American Society of Hematology. 15 November 1998, Vol. 92, No. 10, Supplement 1, Abstract # 668, see abstract.	1-63
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "B" earlier document published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "A" document member of the same patent family		
Date of the actual completion of the international search 12 JUNE 2000		Date of mailing of the international search report 09 AUG 2000
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer KAREN M. HAUDA Telephone No. (703) 308-0196 